



SACRAL NERVE STIMULATION (NEUROMODULATION)

Information about your procedure from
The British Association of Urological Surgeons (BAUS)

This leaflet contains evidence-based information about your proposed urological procedure. We have consulted specialist surgeons during its preparation, so that it represents best practice in UK urology. You should use it in addition to any advice already given to you.

To view the online version of this leaflet, type the text below into your web browser:

[http://www.baus.org.uk/_userfiles/pages/files/Patients/Leaflets/Sacral neuromodulation.pdf](http://www.baus.org.uk/_userfiles/pages/files/Patients/Leaflets/Sacral%20neuromodulation.pdf)

Key Points

- Sacral nerve stimulation (SNS) is used for patients with overactive bladder and for some types of urinary retention in women
- Around 70% of patients are significantly improved after SNS
- SNS involves two separate operations, several weeks apart
- A temporary stimulator is put in first, followed by a two to eight-week “test phase”, and then a second procedure to implant the permanent stimulator
- **The most important complication is wound infection; you should contact your medical team immediately if this occurs**
- You should not have some types of surgical diathermy (electric current used to cut or stop bleeding) after SNS
- You should not go in an MRI scanner (other than for a head scan) after SNS
- You may find that the effect of the stimulator wears off gradually over time as your body gets used to it; re-programming the stimulator may overcome this
- Some patients need a further procedure to change a component of the device if it stops working
- All patients will need further surgery to change the stimulator when the battery runs down (usually around five years)

What does this procedure involve?

Sacral nerve stimulation (SNS) is also sometimes called sacral neuromodulation (SNM). The procedure involves placing a wire in your lower back, near the nerves that control your bladder and bowel. This wire delivers small electrical pulses and helps to improve problems with bladder and bowel function.

It is used to treat the following conditions:

- **Overactive bladder** - frequent and urgent need to pass urine with urine leakage
- Some types of **urinary retention** (inability to empty the bladder) in females
- **Faecal (bowel) incontinence** – and some other bowel problems

For bladder problems, the success rate of SNS is around 70%, meaning that patients report a significant improvement (not cure) in their symptoms. Not all patients are suitable for SNS treatment.

What are the alternatives?

Sacral nerve stimulation may be used to treat two different conditions:

Overactive bladder (OAB)

OAB can be treated without surgery. We recommend that all patients try conservative treatments before having an operation, because it avoids the risks of side-effects or complications of surgery. Treatment options include:

- **Incontinence pads** - if your symptoms are not a bother you may choose to do nothing and use pads for urine leakage
- [Conservative measures](#) – including weight loss, improving fluid intake and reducing caffeine
- [Bladder training](#) – learning techniques to hold on and over-ride your urge to pass urine
- [Medicines](#) – these may help if conservative treatment does not work

Sacral nerve stimulation is only tried if these treatments are not effective. Other procedures that can be used include:

- [Botox injections](#) – botulinum toxin A injected into the wall of your bladder using a telescope
- [Enterocystoplasty](#)– a major operation that enlarges your bladder using a piece of bowel

- **Acupuncture** and **posterior tibial nerve stimulation, PTNS** (electrical stimulation of a nerve near the ankle) - can be used, but are not widely available as an NHS treatment

Urinary retention

If you cannot empty your bladder, SNS sometimes helps but it is not suitable for everyone with this problem. Alternative treatments include:

- **Intermittent self-catheterisation in women** - passing a disposable catheter tube into the bladder through the water pipe to drain the urine
- **Permanent insertion of a catheter**
- **Mitrofanoff procedure** – a major operation to allow you to pass a catheter into the bladder through the abdomen (tummy) instead of the water pipe

What happens on the day of the procedure?


Your surgeon will briefly review your history and medications, and will discuss the surgery again with you to confirm your consent.

Details of the procedure

The treatment involves two separate operations, several weeks apart. After the first part, there is a test phase, which allows the medical team to see if the SNS is working.

During the first operation

- we make a tiny incision (cut) in your lower back and a second 3 - 4 cm incision in your upper buttock
- we place an electrode (wire) near the nerves using X-rays to make sure it is in the correct place; the wire passes out through the skin at the side of your buttock
- we connect this wire to a stimulator box outside your body
- the stimulator box sends electrical signals to the nerves and needs to be worn all the time
- you may be able to home on the same day but an overnight stay is sometimes needed



During the test phase

- You must keep your dressings clean and dry. This means that you cannot shower or bathe during the test phase. You can keep yourself clean by using a sponge or flannel, providing the dressings remain dry. Please do not remove or change the dressings.
- Once you get home, your urology team will monitor you closely. This is so they can check that the treatment is working well, and make any adjustments needed to the stimulator when needed.
- This phase usually lasts between two and eight weeks. It is very important that you complete any [bladder diaries](#) given to you during this time, to see if the treatment is helping your symptoms.

During the second operation

If the SNS treatment does not work during the test phase, we will remove the temporary wire during the second procedure. Otherwise, you will have a permanent stimulator out in.








- we re-open the incision in your buttock and place the permanent stimulator under the skin
- depending on the type of wire put in during the first operation, we either replace it with a new (permanent) wire, or use the same wire
- you may be able to go home on the same day but an overnight stay is sometimes needed
- you may need to take antibiotics to reduce the risk of infection

We switch your stimulator on after the procedure although, sometimes, we leave this for a few days. When it is switched on, you may feel a tapping sensation inside you. We will programme the stimulator so that you get maximum benefit without discomfort.

We will show you how to adjust the stimulator strength yourself, using a remote control.

Are there any after-effects?

The possible after-effects and your risk of getting them are shown below. Some are self-limiting or reversible, but others are not. We have not listed very rare after-effects (occurring in less than 1 in 250 patients) individually. The impact of these after-effects can vary a lot from patient to patient; you should ask your surgeon's advice about the risks and their impact on you as an individual:

After-effect	Risk
Failure of the treatment to improve your symptoms significantly	 More than 1 in 5 patients (over 20%)
Mild discomfort requiring simple painkillers	 Between 1 in 2 & 1 in 10 patients
Need for replacement, relocation or removal of the implanted stimulator or electrode (wire)	 Around 1 in 5 patients (20%)
Discomfort in your buttock at the site of the stimulator	 Around 1 in 10 patients (10%)
Discomfort in your ankle or foot	 Around 1 in 10 patients (10%)
Infection in your wound requiring antibiotics and possible removal of the device	 Between 1 in 10 & 1 in 50 patients
Stimulation produces an adverse (undesirable) after-effect on your bowel function	 Between 1 in 10 & 1 in 50 patients

PLEASE NOTE

Other unwanted side-effects may include unexpected triggering of shop security alarms or airport security scanners. This is not harmful, but you should inform security staff if your device has triggered alarms in the past.

What is my risk of a hospital-acquired infection?

Your risk of getting an infection in hospital is approximately 8 in 100 (8%); this includes getting *MRSA* or a *Clostridium difficile* bowel infection. This figure is higher if you are in a “high-risk” group of patients such as patients who have had:

- long-term drainage tubes (e.g. catheters);
- bladder removal;

- long hospital stays; or
- multiple hospital admissions.

What can I expect when I get home?

- you will get some mild discomfort which may last several days and can normally be relieved with mild painkillers
- you will be given a copy of your discharge summary and a copy will also be sent to your GP
- you must keep the dressings clean and dry
- you must not shower or bathe during the test phase
- you must avoid strenuous activities, sports and any stretching of the lower back for six weeks; these can cause the wire to move out of position so that the SNS stops working
- if you get any flu-like symptoms, redness or throbbing in the wound, pain or burning when passing urine, you should contact the hospital as soon as possible.
- you may need to “fine-tune” your stimulator using the hand-held remote control
- if you do have problems with the stimulator, please contact your named specialist nurse
- a follow-up appointment will be made for you to review your response to the surgery

Are there any long-term issues with sacral nerve stimulation?

Yes, a few. The important ones are:

- **Battery failure** - the battery in your stimulator will run down eventually, usually after four to eight years. Battery life depends on the settings you have been using. Changing the battery is relatively simple, and involves a procedure very like the second stage of the original implantation
- **Electronic devices** - security screening and airport scanning devices can affect your stimulator. Show the security staff your SNS identification card and they may allow you to by-pass the scanner. If not, make sure you turn your simulator off before you pass through the scanner
- **Surgical diathermy** - cutting and coagulating tissue during surgery is usually done using diathermy (electrical current). The current from some types of diathermy equipment can damage your stimulator. Please tell your surgeon about your SNS and he/she will make sure that only safe diathermy equipment is used

- **Sports and strenuous activities** - because of the risk of damaging or dislodging the SNS device or the lead, we advise you not to take part in contact sports, extreme sports or horse-riding
- **Magnetic resonance (MR) scanning** - the powerful magnetic field in an MRI scanner can damage your stimulator and cause you harm. Having an MRI scan of your head will cause no problem, but you should not have any other form of MRI scan

General information about surgical procedures

Before your procedure

Please tell a member of the medical team if you have:

- an implanted foreign body (stent, joint replacement, pacemaker, heart valve, blood vessel graft);
- a regular prescription for a blood thinning agent (warfarin, aspirin, clopidogrel, rivaroxaban or dabigatran);
- a present or previous MRSA infection; or
- a high risk of variant-CJD (e.g. if you have had a corneal transplant, a neurosurgical dural transplant or human growth hormone treatment).

Questions you may wish to ask

If you wish to learn more about what will happen, you can find a list of suggested questions called "[Having An Operation](#)" on the website of the Royal College of Surgeons of England. You may also wish to ask your surgeon for his/her personal results and experience with this procedure.

Before you go home

We will tell you how the procedure went and you should:

- make sure you understand what has been done;
- ask the surgeon if everything went as planned;
- let the staff know if you have any discomfort;
- ask what you can (and cannot) do at home;
- make sure you know what happens next; and
- ask when you can return to normal activities.

We will give you advice about what to look out for when you get home. Your surgeon or nurse will also give you details of who to contact, and how to contact them, in the event of problems.

Smoking and surgery

Ideally, we would prefer you to stop smoking before any procedure. Smoking can worsen some urological conditions and makes complications more likely after surgery. For advice on stopping, you can:

- contact your GP;
- access your local [NHS Smoking Help Online](#); or
- ring the free NHS Smoking Helpline on **0800 169 0 169**.

Driving after surgery

It is your responsibility to make sure you are fit to drive after any surgical procedure. You only need to [contact the DVLA](#) if your ability to drive is likely to be affected for more than three months. If it is, you should check with your insurance company before driving again.

What should I do with this information?

Thank you for taking the trouble to read this information. Please let your urologist (or specialist nurse) know if you would like to have a copy for your own records. If you wish, the medical or nursing staff can also arrange to file a copy in your hospital notes.

What sources have we used to prepare this leaflet?

This leaflet uses information from consensus panels and other evidence-based sources including:

- the [Department of Health \(England\)](#);
- the [Cochrane Collaboration](#); and
- the [National Institute for Health and Care Excellence \(NICE\)](#).

It also follows style guidelines from:

- the [Royal National Institute for Blind People \(RNIB\)](#);
- the [Information Standard](#);
- the [Patient Information Forum](#); and
- the [Plain English Campaign](#).

Disclaimer

We have made every effort to give accurate information but there may still be errors or omissions in this leaflet. BAUS cannot accept responsibility for any loss from action taken (or not taken) as a result of this information.

PLEASE NOTE

The staff at BAUS are not medically trained, and are unable to answer questions about the information provided in this leaflet. If you do have any questions, you should contact your urologist, specialist nurse or GP.